



**Ionization in the Winds of Early-B Stars:  
Constraints Imposed by EUVE**

**J.J. MacFarlane, D.H. Cohen, J.P. Cassinelli, P. Wang**

**June 1995**

**UWFDM-981**

Proceedings of the IAU 152: "Astrophysics in the Extreme Ultraviolet", Berkeley CA,  
27–30 March 1995.

***FUSION TECHNOLOGY INSTITUTE***

***UNIVERSITY OF WISCONSIN***

***MADISON WISCONSIN***

### **DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government, nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

**Ionization in the Winds of Early-B Stars:  
Constraints Imposed by EUVE**

J.J. MacFarlane, D.H. Cohen, J.P. Cassinelli, P.  
Wang

Fusion Technology Institute  
University of Wisconsin  
1500 Engineering Drive  
Madison, WI 53706

<http://fti.neep.wisc.edu>

June 1995

UWFDM-981

# Ionization in the Winds of Early-B Stars: Constraints Imposed by EUVE

By J.J. MAC FARLANE<sup>1,2</sup>, D.H. COHEN<sup>1</sup>  
J.P. CASSINELLI<sup>1</sup> AND P. WANG<sup>2</sup>

<sup>1</sup>Department of Astronomy, University of Wisconsin,  
475 N. Charter St., Madison, WI 53706, USA

<sup>2</sup>Fusion Technology Institute, University of Wisconsin,  
1500 Johnson Drive, Madison, WI 53706, USA

Extreme Ultraviolet Explorer (EUVE) spectral observations of  $\epsilon$  CMa (B2 II) provide significant new information about the ionization dynamics of its wind, its mass loss rate, and the interaction of its wind with the soft X-ray/EUV radiation field. We present results from wind ionization calculations which show how EUV radiation, emitted predominantly in the form of Fe IX–XVI lines from plasma with  $T \sim 1 - 3 \times 10^6$  K, significantly alters the ionization state of its wind. EUV photons from the hot plasma photoionize He II in the cool portion of  $\epsilon$  CMa's wind, producing anomalously high abundances of He III. The subsequent recombination from He III results in He II  $L\alpha$  (304 Å) and  $L\beta$  (256 Å) lines which are observed by EUVE. Also observed is O III  $\lambda\lambda$  374, which results from the O III  $\lambda\lambda$  304 multiplet being pumped by He II  $L\alpha$  — the Bowen mechanism (Cassinelli *et al.* 1995). We report on initial results from numerical simulations which show the effect of the Bowen mechanism on the O III 374 Å line emission.

---

## 1. Introduction

Ultraviolet observations of hot stars (Snow and Morton 1976, Lamers and Morton 1976) have shown that they suffer significant mass loss through high velocity winds. For early-O stars, radio (Bieging, Abbott, and Churchwell 1989) and  $H\alpha$  (Lamers and Leitherer 1993) observations provide evidence for mass loss rates  $\sim 10^{-6} - 10^{-5} M_{\odot}/\text{yr}$ , while wind velocities can be 3,000 km/s or more. Observational determination of mass loss rates for late-O and early-B stars becomes more difficult, however, because their winds are much weaker; i.e., they tend to exhibit lower densities and lower velocities. Fortunately, EUVE observations of the B2 II star  $\epsilon$  CMa (Cassinelli *et al.* 1995) have provided valuable new data for studying the winds of early-B stars. Emission lines between 150 Å and 400 Å have been observed from both the hot X-ray emitting region (Fe IX–XVI) and the cool wind (He II and O III). Below, we present results from wind ionization calculations which show the influence of the soft X-ray/EUV radiation on the ionization state of the wind, and the effect of the Bowen mechanism on the O III 374 Å line intensity.

ROSAT observations of  $\epsilon$  CMa (Drew, Denby, and Hoare 1993) showed that  $\epsilon$  CMa, like most other hot stars, has a significant X-ray luminosity ( $L_x \sim 1 \times 10^{-7} L_{\text{bol}}$ ). Two-temperature fits to the ROSAT data suggest a relatively strong low-temperature component ( $T \sim 1 - 2 \times 10^6$  K), and a weaker high-temperature component ( $T \sim 8 \times 10^6$  K). Plasmas at these temperatures also exhibit strong Fe line emission between 150 Å and 400 Å, which was observed by EUVE for  $\epsilon$  CMa. Recent studies (MacFarlane, Cohen, and Wang 1994) have predicted that the soft X-ray/EUV radiation of early-B stars can significantly alter the bulk ionization state of their winds. This is unlike the case of early-O star winds, where X-rays tend to produce only a small (but detectable) perturbation on their ionization distribution (Cassinelli and Olson 1979).

Because EUVE has detected both the hot plasma Fe line emission and cool wind He II

and O III line emission, it presents a rare opportunity to study the interplay of the hot plasma radiation field and the cool wind.

In this paper, we present several results from numerical simulations describing these effects. More detailed descriptions of this work will be presented elsewhere (Cohen *et al.* 1995, MacFarlane *et al.* 1995).

## 2. Models

Detailed descriptions of our wind ionization code have been presented elsewhere (MacFarlane *et al.* 1993, 1994); only a brief summary is presented here. The wind is assumed to be spherically symmetric with a velocity which increases monotonically with radius. The density is specified by its mass loss rate and assuming, the mass flux is constant with radius. The radiation field includes contributions from: (1) the photosphere, based on Mihalas' (1972) non-LTE models longward of 912 Å and a fit to the EUVE data for 350 Å < λ < 912 Å; (2) diffuse radiation in the wind; and (3) EUV/X-ray radiation from a high-temperature plasma, which had a frequency-dependence given by XSPEC models (Raymond and Smith 1977) and a power-law temperature distribution determined from the combined analysis of ROSAT and EUVE data (Cohen *et al.* 1995). Emission from the hot plasma was assumed to be distributed throughout the wind (shock model), with a peak in the differential emission measure at  $r = 1.5 R_*$  ( $\alpha = -3$  in Eqn. (10) of MacFarlane *et al.* 1993). Radiative transfer effects for all radiation sources were computed using a multi-ray impact parameter model.

Non-LTE atomic level populations were computed by solving multilevel statistical equilibrium equations self-consistently with the radiation field. A total of 81 levels for H, He, and O were considered in our atomic model. It was necessary to consider fine-structure splitting of the O III levels so that the Bowen photoexcitation mechanism (Bowen 1935) could be accounted for. This mechanism arises due to overlap of the He II  $L\alpha$  line at 303.78 Å and the O III  $1s^2 2s^2 2p^2 \ ^3P - 1s^2 2s^2 2p 3d \ ^3P^0$  multiplet (see Figure 1). When He III recombines to the  $n = 2$  state of He II, the  $L\alpha$  photons emitted can be absorbed, producing an anomalously high population of the O III  $3d \ ^3P^0$  level. This level subsequently decays to the  $3s \ ^3P^0$  level, resulting in the spontaneous emission of the  $\lambda\lambda$  374 multiplet, which is observed as a single emission feature by EUVE (Cassinelli *et al.* 1995). The interaction of overlapping lines is modeled using a generalized Sobolev method with multiple velocity surfaces (Rybicki and Hummer 1978; Puls, Owocki, and Fullerton 1993; see also MacFarlane *et al.* 1995).

## 3. Results

We first examine the effect of soft X-ray/EUV radiation on the He ionization balance in the wind of  $\epsilon$  CMa. Figure 2 shows computed ionization distributions for He I-III from two calculations: one using a distributed X-ray emission source (solid curves) and the other with no X-ray source (dotted curves). In each case the mass loss rate was  $0.8 \times 10^{-8} M_{\odot}/\text{yr}$ . Clearly, the population of doubly ionized He (circles) is dominated by photoionization due to the hot plasma radiation, which dominates the radiation field at  $h\nu > 54$  eV. At relatively low velocities ( $v \leq 0.4 v_{\infty}$  and  $r < 1.5 R_*$ ) the He III population is lower because of the higher wind densities and the fact that the X-ray differential emission measure peaks at  $r = 1.5 R_*$  in this model. Note that in the absence of X-ray-induced photoionization, the He III fraction is  $\leq 10^{-4}$  throughout the wind. The distribution of He III in the wind is important because its recombination is the source of photons which are ultimately observed by EUVE at 304 Å and 374 Å.




FIGURE 1. Schematic energy level diagram illustrating radiative processes in the wind of  $\epsilon$  CMa.




FIGURE 2. Helium ionization distributions calculated with a distributed hot plasma radiation source (solid curves) and with no X-ray radiation source (dashed curves).




FIGURE 3. Calculated O III  $\lambda\lambda$  374 line profiles for  $\epsilon$  CMa wind ionization models with  $\dot{M} = 0.8 \times 10^{-8} M_{\odot}/\text{yr}$ . Dashed curve: atomic populations computed without line overlap effects; solid curve: with line overlap effects.

The effect of the Bowen mechanism (photopumping of the O III  $\lambda$  304 multiplet by He II  $\text{L}\alpha$ ) on the O III  $\lambda\lambda$  374 flux can be seen in Figure 3, where the 374 Å profile is shown from two calculations. The solid curve shows the profile from a calculation in which line overlap effects on the atomic populations were included, while the dashed curve is from a calculation in which photoexcitation due to overlapping lines was neglected. (Note, however, that the emergent spectra, which are computed after the level populations are determined, *do* include overlap effects. This gives rise to the structure seen in the absorption component of the profile.) By including the Bowen effect in the calculation, the emission from the 374 Å multiplet is seen to increase substantially. Without the Bowen effect, the  $\lambda\lambda$  374 profile has the characteristic P-Cygni shape, with blue-shifted absorption and red-shifted emission; that is, scattered photospheric radiation with no additional source of photons. Preliminary results suggest that the best agreement with the EUVE data for the 304 Å and 374 Å lines is obtained for mass loss rates of  $\sim 1 - 2 \times 10^{-8} M_{\odot}/\text{yr}$ . Final results and a complete description of this analysis will be presented in a forthcoming paper.

This work has been supported in part by NASA grants NAGW-2210 and NAGW 5-2282.

## REFERENCES

- BIEGING, J. H., ABBOTT, D. C. & CHURCHWELL, E. 1989 *Ap. J.* **340**, 518.
- BOWEN, I. S. 1935 *Ap. J.* **81**, 1.
- CASSINELLI, J. P. & OLSON, R. E. 1979 *Ap. J.* **229**, 304.
- CASSINELLI, J. P., COHEN, D. H., MACFARLANE, J. J., DREW, J. E., LYNAS-GRAY, A. E., HOARE, M. G., VALLERGA, J. V., WELSH, B. Y., VEDDER, P. W., HUBENY, I., & LANZ, T. 1995 *Ap. J.* **438**, 932.
- COHEN, D. H. *et al.* 1995 to be submitted to *Ap. J.*
- DREW, J. E., DENBY, M. & HOARE, M. G. 1994 *MNRAS* **266**, 917.
- LAMERS, H. J. G. L. M. & MORTON, D. C. 1976 *Ap. J. Suppl.* **32**, 715.
- LAMERS, H. J. G. L. M. & LEITHERER, C. 1993 *Ap. J.* **412**, 771.
- MACFARLANE, J. J., COHEN, D. H., & WANG, P. 1994 *Ap. J.* **437**, 351.
- MACFARLANE, J. J., WALDRON, W. L., CORCORAN, M. F., WOLFF, M. J., WANG, P., & CASSINELLI, J. P. 1993 *Ap. J.* **419**, 813.
- MACFARLANE, J. J., COHEN, D. H., CASSINELLI, J. P. & WANG, P. 1995 to be submitted to *Ap. J.*
- MIHALAS, D. 1972 *NCAR Tech. Note*, STR-76.
- PULS, J., OWOCKI, S. P., AND FULLERTON, A. W. 1993 *Astron. Astrophys.* **279**, 457.
- RAYMOND, J. C. & SMITH, B. W. 1977 *Ap. J. Suppl.* **35**, 419.
- RYBICKI, G. B. & HUMMER, D. G. 1978 *Ap. J.* **219**, 654.
- SNOW, T. P. & MORTON, D. C. 1976 *Ap. J. Suppl.* **32**, 429.